


DOCUMENT SCAN CONTROL SHEET

Document ID 505805



EPA # 961-362

EPA #
For SLN

Product LEBANON PROSCAPE FERTILIZER WITH BARRICADE 0.43%
Name PREEMERGENCE WEED CONTROL

Accepted
Date 3/13/2006

Document
Type PRILAB

Registration ID

LEBANON
ProScape FERTILIZER WITH **BARRICADE**® .43%
 PREEMERGENCE WEED CONTROL

SPECIMEN LABEL

LEBANON
ProScape®

BARRICADE® .43%
 PREEMERGENCE WEED CONTROL
 WITH
LEBANON
ProScape FERTILIZER

FERTILIZER
 WITH
BARRICADE® .43%
 PREEMERGENCE WEED CONTROL
16-4-8
MESA

GUARANTEED ANALYSIS

16% Nitrogen (N)	4.3% BARRICADE (N)
4.3% Ammoniacal Nitrogen	
1.2% Urea Nitrogen	
2.8% Urea Nitrogen	
2.2% Urea Nitrogen	
Ammoniacal Phosphate (P ₂ O ₅)	4%
Super Phosphate (P ₂ O ₅)	0%
0.8% Potassium (K ₂ O)	2.8%
2.8% Combined Sulfur (S)	
Residue Sulfur: Ammonium Phosphate, Ammonium Sulfate, Methylene Urea, Urea, Molar of Residue	
Chloride (Cl) as max. Res.	7.0%
27% Shaded Ammoniacal Nitrogen has Methylene Urea	10%

22-53614

SPREADER SETTINGS	Rate in pounds per 1,000 Sq. Ft.	These settings apply to all turfgrass and lawn applications. However, type and condition of substrate, source of material, and amount of water may require slightly different settings for different coverage.
ProScape® 16-4-8 MESA (N-P-K)	3.2	
ProScape® 16-4-8 MESA (N-P-K) (N)	N	
ProScape® 16-4-8 MESA (N-P-K) (P)	P	
ProScape® 16-4-8 MESA (N-P-K) (K)	K	
ProScape® 16-4-8 MESA (N-P-K) (S)	S	

For selective preemergence control of grass and broadleaf weeds in:

- established turfgrasses (excluding golf course putting greens) and lawns
- landscape ornamentals (including established perennials & wildflower plantings)

ACTIVE INGREDIENT:
 Proflarazin: (N,N-Di-[1-propyl-2,4-dinitro-5-(trifluoromethyl)-1H]phenyl)benzamide 0.43%
INERT INGREDIENTS: 99.57%
TOTAL: 100.00%

EPK Reg. No. 561-242 EPA Est. 99-196-01-242-41-01
 Barricade is a Registered Trademark of a Syngenta Group Company Syngentatraz is a first letter of lot number

READ THE ENTIRE LABEL BEFORE USING THIS PRODUCT. USE ONLY ACCORDING TO LABEL INSTRUCTIONS.
 Before buying this product, read the Warranty Section elsewhere on this label.
 If terms are not acceptable, return the unopened container to the place of purchase at once.

KEEP OUT OF REACH OF CHILDREN
CAUTION
 See the following section: Precautionary Statements, Hazards to Humans and Domestic Animals, for additional information.

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION

Causes moderate eye irritation if dusted. Irritation if inhaled or absorbed through the skin. Avoid contact with skin, eyes, or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling. Prolonged or repeated contact may irritate while handling the material may cause dizziness or headache in some individuals.

FIRST AID
 IF IN EYES: Hold eye open and flush slowly and gently with water for 15-20 minutes.
 • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
 • Call a physician or contact poison control for further treatment advice.
 IF ON SKIN: Wash thoroughly with soap and water.
 IF ON CLOTHING: Remove contaminated clothing and wash separately before reuse.
 IF INHALED: Move person to fresh air.
 • If breathing is difficult, call 911 or an ambulance. Do not give artificial respiration, unless directed to do so.
 • Call a physician or contact poison control for further treatment advice.
 Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-235-6236 for emergency medical treatment information.

ENVIRONMENTAL HAZARDS
 This product has low toxicity to water, fish, birds, and bees. The product is not toxic to fish. However, at concentrations above the level of water solubility, it may be toxic to fish. Do not apply directly to water, to areas where surface water is present, or to riparian areas below the means of water flow. Do not apply directly to streams, rivers, or other bodies of water. Do not apply to aquatic organisms or adjacent sites. Do not contaminate water when disposing of equipment and containers.
 If material is spilled: Call a professional service or manufacturer for disposal or use per label instructions. Wash clothes separately to prevent contamination. Do not use clothing that has been contaminated with this product. Wash clothes separately to prevent contamination. Do not use clothing that has been contaminated with this product. Wash clothes separately to prevent contamination. Do not use clothing that has been contaminated with this product.

ACCEPTED FOR REGISTRATION

MAR 13 2006

NEW YORK STATE DEPARTMENT OF ENVIRONMENTAL CONSERVATION
 NET WEIGHT **50** LBS. (22.68 KG)

Dec 1d 505805

FERTILIZER

WITH

BARRICADE® .43%

PREEMERGENCE WEED CONTROL

16-4-8

MESA

P A T E N T E D

GUARANTEED ANALYSIS

Total Nitrogen (N)*	16%
4.0% Ammoniacal Nitrogen	
1.2% Water Insoluble Nitrogen	
8.3% Urea Nitrogen	
2.5% Other Water Soluble Nitrogen	
Available Phosphate (P ₂ O ₅)	4%
Soluble Potash (K ₂ O)	8%
Total Sulfur (S)	2.8%
2.8% Combined Sulfur (S)	

Nutrient Sources: Ammonium Phosphate, Ammonium Sulfate, Methylene Ureas, Urea, Muriate of Potash.

Chlorine (Cl) not more than	7.0%
*3.7% Slowly Available Nitrogen from Methylene Ureas.	F699

MESA®, PennMulch® and ProScape® are registered trademarks of Lebanon Seaboard Corporation.



22-53614

SPREADER SETTINGS	Rate in pounds per 1,000 Sq. Ft.	
	5.3	
ProScape® SS/PrizeLAWN® BF-1/SS/CBR III	N	
PennMulch® HVO/PrizeLAWN® BF-HVO	N	
Earthway Rotary	18	
Lesco (letter dial/numeric dial)	K / 22	
Scotts AccuPro 2000/SR-2000	O	
Scotts R8A Rotary	N	
Scotts R7X Rotary	J	
Spyker Rotary	5	

These settings were calibrated and field tested. However, age and condition of spreader, speed of operation, and evenness of terrain may require slightly different settings for desired coverage.

For selective preemergence control of grass and broadleaf weeds in:

- established turfgrasses (excluding golf course putting greens) and lawns
- landscape ornamentals (including established perennials & wildflower plantings)

SPECIMEN LABEL

ACTIVE INGREDIENT:

Prodiamine: [N³,N³-Di-n-propyl-2,4-dinitro-6-(trifluoromethyl)-
m-phenylenediamine 0.43%

INERT INGREDIENTS: 99.57%

TOTAL 100.00%

EPA Reg. No. 961-362

Barricade® is a Registered Trademark of a Syngenta Group Company.

(L) (D)

EPA Est. 961-PA-01:961-IL-01

Superscript used is first letter of lot number.

READ THE ENTIRE LABEL BEFORE USING THIS PRODUCT. USE ONLY ACCORDING TO LABEL INSTRUCTIONS.

Before buying this product, read the Warranty Section elsewhere on this label.

If terms are not acceptable, return the unopened container to the place of purchase at once.

KEEP OUT OF REACH OF CHILDREN

CAUTION

See the following section, **Precautionary Statements, Hazards To Humans and Domestic Animals**, for additional information.

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION**

Causes moderate eye injury (irritation). Harmful if inhaled or absorbed through the skin. Avoid contact with skin, eyes, or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling. Prolonged or frequently repeated skin contact while handling the material may cause allergic reactions in some individuals.

FIRST AID

IF IN EYES: • Hold eye open and rinse slowly and gently with water for 15-20 minutes.
• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
• Call a poison control center or doctor for treatment advice.

IF ON SKIN • Take off contaminated clothing.

OR CLOTHING: • Rinse skin immediately with plenty of water for 15-20 minutes.
• Call a poison control center or doctor for treatment advice.

IF INHALED: • Move person to fresh air.
• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.
• Call a poison control center or doctor for further treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact (888) 208-1368 for emergency medical treatment information.

ENVIRONMENTAL HAZARDS

This product has low solubility in water. At the limits of solubility, this product is not toxic to fish. However, at concentrations above the level of water solubility, it may be toxic to fish. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff from treated areas may be hazardous to aquatic organisms in adjacent sites. Do not contaminate water when disposing of equipment washwaters.

If material is spilled: Contain and sweep up material of spill for disposal or use per label instructions. Wash cleanup equipment to prevent contamination elsewhere. Excess quantities above normal application rates could result in damage to plants subject to spill. Water-in remainder, discard under, cover with layer of soil, or leave exposed to sunlight, as deemed appropriate to the situation.

Lebanon Seaboard Corporation

1600 E. Cumberland St. • Lebanon, PA 17042

www.LebanonTurf.com

3 HT 50 LBS. (22.68 KG)

SPECIMEN LABEL

LEBANON ProScape®

FERTILIZER WITH BARRICADE® .43% PREEMERGENCE WEED CONTROL

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

STORAGE: Store this product only in its original container in a dry, cool, secured storage area, away from other fertilizer, feed, or food stuffs and separated from other pesticides.

DISPOSAL: If empty: Do not reuse this container. Place in trash or offer for recycling if available.

If partly filled: Call your local solid waste agency or 1-800 CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

This product is a selective preemergence herbicide that provides residual control of many grass and broadleaf weeds in established turfgrasses (excluding golf course putting greens) and lawns, landscape ornamentals, established perennials and wildflower plantings.

This product controls susceptible weeds by inhibiting weed seeds germination and root development. Most effective weed control will be obtained when this product is activated by at least 0.5 inch of rainfall or irrigation, or with shallow (1-2 inches) incorporation, prior to weed seed germination and within 14 days following application.

Not for use on plants being grown (1) for sale or other commercial use, (2) for commercial seed production, or (3) for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in ornamental gardens or parks, or on golf courses or lawns and grounds.

Do not graze or feed livestock forage cut from areas treated with this product.

Do not apply this product through any type of irrigation system.

Do not apply aerially.

Do not apply to golf course putting greens.

APPLICATION DIRECTIONS

Apply uniformly with suitable, calibrated application equipment.

ESTABLISHED TURF

This product is a selective preemergence herbicide that, when properly applied, will control certain grass and broadleaf weeds in established turfgrasses and lawns. The maximum amount of this product that may be applied per year is given for each turfgrass species in the **Maximum Annual Rates** section of this label.

Most effective weed control in turfgrasses will be obtained when this product is activated by at least 0.5 inch of rainfall or irrigation prior to weed seed germination and within 14 days following application. See the map below for approximate crabgrass seed germination dates.

CRABGRASS SEED GERMINATION DATES



Approximate Date
After May 30
After May 10
After April 20
After March 20
January 1 to March 20

USE PRECAUTIONS

The following precautions apply to the use of this product in turfgrasses and lawns:

Application of this product may thin emerged annual bluegrass and newly overseeded grasses.

Do not apply to overseeded turf within 60 days after seeding or until after the second mowing, whichever is longer. Injury to desirable seedlings is likely if this product is applied before seeding secondary roots are in the second inch of soil, not thatch plus soil.

Do not cut (harvest) treated sod before 120 days after application. Do not apply to newly set sod until the following year.

Application of this product to turf stressed by drought, low fertility, or pest damage may result in turf injury.

Disturbing the herbicide barrier with cultural practices such as disking may result in reduced weed control.

Do not apply this product to putting greens or areas where dichondra, colonial bentgrass, velvet bentgrass or annual bluegrass (*Poa annua*) are desirable species.

RATES OF APPLICATION

This product may be applied as a single application or in sequential applications to control weeds germinating throughout the year. All applications must be made prior to germination of the target weeds. This product will not control established weeds.

Maximum use rate selection should be based on turf species. The length of time of residual weed control provided by this product is related to the rate applied.

MAXIMUM ANNUAL RATES

This product is recommended for use on the turfgrass species listed in the following table. Do not exceed the maximum yearly rate as given in the following table:

MAXIMUM APPLICATION RATE/CALENDAR YEAR of LEBANON PROSCAPE FERTILIZER WITH BARRICADE .43% PREEMERGENCE WEED CONTROL by Turfgrass Species¹

Turf Species:	Lbs. Product/Acre	Lbs. AI/Acre	Lbs./1,000 Sq. Ft.
Bermudagrass ² , Bahiagrass, Centipedegrass, Seashore Paspalum, St. Augustinegrass, Tall Fescue (including turf-type), Zoysia	349	1.5	8.0
Buffalograss, Kentucky Bluegrass, Perennial Ryegrass	233	1.0	5.3
Creeping Red Fescue	175	0.75	4.0
Creeping Bentgrass	150	0.65	3.4

STANDARD GREEN LABEL

¹⁾ These are the maximum rates per calendar year by species limitations.

²⁾ May be used on newly sprigged or plugged Bermudagrass at rates not to exceed 121 lbs./Acre. Newly sprigged or plugged Bermudagrass stolon rooting may be temporarily retarded. Suppression only of Foxtail, Goosegrass and Rescuegrass due to reduced product rates used in sprigging situations.

• Do not apply more than 349 lbs. of this product per calendar year.

• Use higher rates of this product to achieve higher levels of fertility and longer periods of weed control for each turf type, but do not exceed the maximum application rates specified in the **Maximum Annual Rates Table**.

WEEDS CONTROLLED

When used in accordance with this label this product will provide control of the following weeds:

Barnyardgrass	Chickweed, Mouseear (from seed)	Foxtails, Annual	Johnsongrass (from seed)	Lambsquarter, Common	Purslane, Common	Speedwell, Persian
Bluegrass, Annual (<i>Poa annua</i>)	Crabgrass (large, smooth)	Goosegrass ¹	Junglerice	Lovegrass	Pusley, Florida	Sprangletop
Carpetweed	Crowfootgrass	Henbit	Knotweed	Panicum (Texas, Fall, Browntop)	Rescuegrass ²	Spurge, Prostrate
Chickweed, Common	Cupgrass, Woolly	Itchgrass	Kochia	Pigweed	Shepherd's Purse ²	Witchgrass
					Signalgrass, Broadleaf	Woodsorrel, Yellow (from seed)

¹ In many areas a single application of 150 to 349 lbs./Acre of this product will control goosegrass. However, under heavy goosegrass pressure and/or an extended growing season, most effective weed control may be obtained by making an initial application of 150 to 233 lbs./Acre followed after 60 to 90 days by a second application at doses that would not exceed those given in the **Maximum Annual Rate Table**. Do not exceed the maximum rate for turfgrass species listed in the **Maximum Annual Rate Table** above.

² Applications for this weed should be made in late summer, fall or winter prior to germination.

³ Suppression only.

Sequential applications may be made so long as the total amount of product applied does not exceed the maximum annual application rates recommended for each turf species. All applications must be made prior to germination of the weed seeds.

WHEN TO APPLY AFTER OVERSEEDING TURF

Do not apply to overseeded turf within 60 days after seeding or until after the second mowing, whichever is longer. Injury to desirable seedlings is likely if this product is applied before seeding secondary roots are in the second inch of soil, not thatch plus soil.

WHEN TO OVERSEED AFTER APPLICATION

This product will inhibit the germination of turf species if overseeded too soon after application. Follow rates and intervals in the table below for best overseeding/reseeding results.

LEBANON PROSCAPE FERTILIZER WITH BARRICADE .43% PREEMERGENCE WEED CONTROL Rate (lb./A)		Months Before Overseeding		
Product	AI	North	Transition	South
116	.50	4	4	4
150	.66	5	4	4
175	.74	6	5	5
185	.80	---	6	6
233	1.00	---	7	7
265	1.14	---	---	9
302	1.30	---	---	10
349	1.50	---	---	12

LANDSCAPE ORNAMENTALS (including Established Perennials and Wildflower Plantings)

This product may be applied for residual preemergence weed control in ornamentals.

USE RATES

Apply this product at 150 to 349 lbs./Acre in fall and/or spring. Use higher rates of application for longer control periods. Sequential applications may be made so long as the total amount of product applied does not exceed the maximum annual application rate of 349 lbs./Acre.

APPLICATION TIMING AND INFORMATION

This product is a preemergence herbicide and will not control emerged weeds. Most effective weed control in ornamentals will be obtained when this product is activated by at least 0.5 inch of rainfall or irrigation, or with shallow (1 to 2 inches) mechanical incorporation, prior to weed seed germination and within 14 days following application.

Best weed control is obtained when this product is applied to soil free of clods, weeds, and debris such as leaves. Prior to application of this product, control existing vegetation by hand weeding, cultivation or the use of an appropriate postemergence herbicide.

ESTABLISHED LANDSCAPE ORNAMENTALS

This product may be applied at any time to established ornamentals as a broadcast, over the top or directed application. Irrigation or rainfall soon after application will wash residues off plant foliage and activate this product in the soil.

NEWLY TRANSPLANTED LANDSCAPE ORNAMENTALS

This product may be applied to newly transplanted ornamentals as a broadcast, over the top or directed application. Delay application to allow soil to settle around new transplants and water thoroughly before applying this product. Apply this product after cuttings form roots and become established. Apply before budding/grafting or after buds/grfts have taken to avoid any inhibition of the tissue union.

TOLERANT ORNAMENTAL SPECIES

This product will not harm most trees, shrubs, vines and flowers. The species listed below are tolerant to this product. Best results will be obtained when this product is uniformly applied to the soil surface; avoid applying to ornamental shoots which may defeat the effort to get uniform distribution on the soil surface. This product may be applied over the top of these species.

When plants are under stress (such as heat, drought, or frost damage) some cultivars of listed plants may be sensitive to this product.

Fir species** (Balsam, Fraser, Noble, etc.)	Feathery Cassia Wild Lilac	<i>Gaura</i> sp. Gentian	Edelweiss Chinese Privet; Variegata	Canary Island Pine Slash Pine	'White Gumbo' Lady Bank's Rose
Abelia; Sherwood	<i>Ceratostigma plumbagonoides</i>	Cranesbill	Amur Privet	Aleppo Pine	Rosemary*
Japanese Maple	False Cypress	Gladiolus species**	Japanese Privet	Austrian Pine	Black-Eyed Susan; Goldstrum
Norway Maple	<i>Chrysanthemum nipponicum</i>	Baby's Breath	Glossy Privet (Waxleaf)	Longleaf Pine	Santolina
Yarrow; King Edward	Cleyera	English Ivy	Lily; Jazz	Monterey Pine	Saxifrage; Purple Dome
Kiwi*	Citrus species*	Sunrose	Big Blue Lily; Turf; Silver Mound;	Japanese Black Pine	Pincushion Flower
Lily-of-the-Nile (African Lily)	Coreopsis (Calliopsis): Early	Daylily; Aztec Gold; Stella De Oro,	Evergreen Giant	Eastern White Pine	Stoncrop (<i>Sedum album</i>)
<i>Agapanthus orientalis</i>	Sunrise, Moonbeam	Tender Love	Liriope, Variegated	Scotch Pine	Stoncrop; Lidakense
Five-Leaf or Chocolate Vine	Flowering Dogwood	Rose of Sharon**	Liriope, Creeping	Loblolly Pine	Stoncrop (<i>Sedum dasycyllum</i>)
Lady's Leek, Nodding Onion	American Dogwood	Mallow; Disco Belle White	Cardinal Flower, Indian Pink	Virginia Pine	Stoncrop; Dragon's Blood
Japanese Anemone	Pampas Grass	Chinese Hibiscus	Japanese Honeysuckle	Pistachio*	Spirea; Anthony Waterer
Aquilegia; Red and Gold	Cranberry Cotoneaster	Hosta, Plantain Lily (Fragrant)	Tatarian Honeysuckle	Queensland Pittosporum	Australian Brushcherry
Vine Hill Manzanita	Cotoneaster	Hosta, "Searsucker"	Burgundy	Japanese Pittosporum	Japanese Yew
Cape Weed	Bearberry Cotoneaster	<i>Houttuynia cordata</i> var. <i>variegata</i>	Loosestrife; Modern Pink	Japanese Yew	Yew
Wormwood; Silver Mound, Castle	Rockspray Cotoneaster	Holly	Magnolia species**	English Laurel	Germander
Aster; Bonny Blue, Purple Dome	Hawthorne	Chinese Holly	Ice Plant	Almond, Apricot, Nectarine, Peach,	Meadow Rue
Aster <i>X Frikartii</i>	Lucifer	Japanese Holly; Helleri	Crabapple*	Plum and Prune*	American Arborvitae
Lady Fern; Fern Lady	Italian Cypress	American Holly	Yaku Jima, Silberfeder	Douglas Fir**	Star Jasmine
Japanese Aucuba	White Trailing Ice Plant	Yaupon Holly; Schillings	Heavenly Bamboo	Firethorn (Scarlet)	Canada Hemlock
Fibrous Begonia; Hardy Grandis	Cooperi Pink	<i>Inula ensifolia</i>	Narcissus species**	Firethorn (<i>P. Fortuneana</i>)	Tulip species
Barberry	Larkspur; Blue Elf	Sword-Leaved Iris; Jodlesong	Oleander	Firethorn (<i>P. Koidzumii</i>)	Veronica, Speedwell; Sunny Border
Wintergreen Barberry	Dianthus; Maiden Pinks "Zing"	Siberian Iris; Cabemet	Olive*	Bradford Pear sp.	Japanese Viburnum
Mentor; Barberry	Cheddar Pink	Iris species**	Mondo Grass	Oak, Shumard's Red	Sweet Viburnum
Japanese Barberry	Hop Bush	Winter Jasmine	Osmanthus (False Holly); Gulf Tide	Oak species	Japanese Snowball
Warty Barberry	Coneflower, Purple; Magnus	Chinese Juniper; Nick's Compact,	Trailing African Daisy	Indian Hawthorne	Canary Island Viburnum
<i>Bergenia cordifolia</i>	Silverberry	Parsonii	Sourwood	Yedda Hawthorn	Arrowood Viburnum
Snowbank	Wintercreeper	Shore Juniper; Blue Pacific	Tree Peony	'Coral Bells'	Laurustinus
Butterfly-Bush (Dwarf Blue);	Japanese Spindle Tree (Evergreen	Parsoni	Fontain Grass (Dwarf)	'Delaware Valley White'	Cranberry Bush
Royal Red	Euonymus)	Creeping Juniper	<i>Perovskia atriplicifolia</i>	'Flame Creeper'	Leatherleaf Viburnum
Japanese Boxwood	Spreading Euonymus	Walnut*	Avocado*	'Formosa'	Greater Periwinkle, Vinca
Weeping Bottlebrush	Japanese Aralia	Shrimp Plant	Frasier's Photinia (Redtip)	'Girard Crimson'	Common Periwinkle, Vinca
Scotch Heather	Border Forsythia		Dragonhead, False; Vivid	'George L. Tabor'	Grape*

Tussock Bellflower, (White Clips)
Trumpet Creeper, Trumpet Flower,
Madame Galen
Hottentot Fig (Ice Plant)

Weeping Forsythia
Greenstem Forsythia
Gaillardia, Blanket Flower, "Goblin"
Gardenia, Cape-Jasmine, Radicans

Crape Myrtle
Crape Myrtle; Tuscarora
Weeping Lantana
Lavender; Munstead

Spruce species** (Colorado Blue,
Norway, etc.)
Lily-of-the-Valley Bush
Calabrian Pine

'Hino-crimson'
'PJM'
'Roseum Elegans'
'Wakelebisu'

Old Fashioned Weigela
Spanish Bayonet, Yucca
Adam's Needle, Yucca

* Do not use on food producing trees, vines, or plants. ** Not for use on container grown plants.

NOTE: Supplemental applications of lawn fertilizer may be needed to achieve a minimum of 1 pound nitrogen per 1,000 sq. ft.

LIMITATION OF WARRANTY AND LIMITATION OF LIABILITY

NOTICE: Read this Limitation of Warranty and Limitation of Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of The Manufacturer or Seller. All such risks shall be assumed by Buyer or User.

The Manufacturer warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, under normal use conditions, subject to the risks described above. THE MANUFACTURER MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR OF MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY. In no event shall The Manufacturer or Seller be liable for any incidental, consequential or special damages resulting from the use or handling of this product.

The Manufacturer and Seller offer this product, and Buyer and User accept it, subject to the foregoing limitations of warranty and limitation of liability, which may not be modified by any oral or written agreement.

MANUFACTURED BY:
LEBANON SEABOARD CORPORATION
1600 EAST CUMBERLAND STREET
LEBANON, PA 17042
www.LebanonTurf.com

Information regarding the contents and levels of metals in this product is available
on the Internet at <http://www.regulatory-info-lebsea.com>

SPECIMEN LABEL