

DOCUMENT SCAN CONTROL SHEET

Document ID 513712

EPA # 79676-21

EPA #
For SLN

Product LESCO SPECTATOR TURF AND ORNAMENTAL FUNGICIDE
Name

Accepted
Date 1/30/2008

Document
Type PRILAB

Registration ID 125954

e-1

Doc ID: 513712

ACCEPTED FOR REGISTRATION

JAN 30 2008

NYSDEC

LESCO[®] SPECTATOR[™] Turf and Ornamental Fungicide

#450267
NET CONTENTS: 1 qt

For use on turf and ornamentals and non-bearing fruit and nut trees in nursery and landscape plantings

ACTIVE INGREDIENT:

Propiconazole: 1-[[2-(2,4-dichlorophenyl)-4-propyl-1,3-dioxolan-2-yl]methyl]-1H-1,2,4-triazole 41.8%

OTHER INGREDIENTS: 58.2%

TOTAL: 100.0%

Contains petroleum distillates. Contains 4 lbs. of propiconazole per gallon.

KEEP OUT OF REACH OF CHILDREN WARNING AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand this label, find someone to explain it to you in detail).

See inside label booklet for First Aid and Precautionary Statements.

EPA Reg. No. 79676-21 EPA Est. No. indicated by the 8th digit of the batch number on this package. (A) = 4-NY-001; (P) = 34704-MS-002
Manufactured for: Etigra • 501 Cascade Pointe Lane, Suite 103, Cary, NC 27513 • www.etigra.com

FIRST AID	
If in eyes:	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15-20 minutes.• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.• Call a poison control center or doctor for treatment advice.
If on skin or clothing:	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15-20 minutes.• Call a poison control center or doctor for treatment advice.
If swallowed:	<ul style="list-style-type: none">• Call poison control center or doctor immediately for treatment advice.• Have person sip a glass of water if able to swallow.• Do not induce vomiting unless told to do so by the poison control center or doctor.• Do not give anything by mouth to an unconscious person.
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-424-9300 for emergency medical treatment information.	
NOTE TO PHYSICIAN	
There is no specific antidote for this product. Induce emesis or lavage stomach, taking care to avoid aspiration of stomach contents into lungs. Give saline laxative and supportive therapy. Contains petroleum distillate – vomiting may cause aspiration pneumonia.	

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

WARNING

Causes substantial but temporary eye injury. Causes skin irritation. Harmful if swallowed. Do not get in eyes, on skin, or on clothing. Wear protective eyewear such as goggles, face shield, or safety glasses. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove and wash contaminated clothing before reuse.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category G on the EPA chemical resistance category selection chart.

Applicators and other handlers must wear:

- Coveralls over short-sleeved shirt and long pants
- Chemical resistant gloves, such as a barrier laminate or viton thicker than 14 mils
- Chemical-resistant footwear plus socks
- Goggles or face shield
- Chemical-resistant headgear for overhead exposures.

Mixers, loaders and persons cleaning equipment must wear:

- Chemical-resistant apron.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Control Statements

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40CFR 160.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

PHYSICAL OR CHEMICAL HAZARDS

Combustible. Do not use or store near heat or open flame.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

(continued)

AGRICULTURAL USE REQUIREMENTS (continued)

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours. Exception: If the product is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water is:

- Coveralls
- Chemical-resistant gloves, such as barrier laminate or viton
- Shoes plus socks
- Goggles or face shield

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard (WPS) for Agricultural Pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Do not enter into treated areas or allow others to enter treated areas without protective clothing until sprays have dried.

GENERAL INFORMATION

Read all sections of this label before using this product. LESCO® Spectator™ Turf and Ornamental Fungicide is a systemic fungicide that controls numerous diseases on turf-grasses, ornamentals, nursery plantings, and landscape plantings; please refer to the appropriate section for the diseases controlled.

USE PRECAUTIONS/RESTRICTIONS

- To prevent plant injury or lack of efficacy, follow all directions for use and precautions on this label.

- Do not apply this product through any type of irrigation system.
- Do not use this product as a tree injection treatment.
- Bermudagrass may be sensitive to LESCO® Spectator™ Turf and Ornamental Fungicide. Do not exceed 1.44 fl. oz. per 1000 sq. ft. per 30 days on any variety of bermudagrass.
- In Florida, do not apply LESCO® Spectator™ Turf and Ornamental Fungicide to bermudagrass golf course greens when temperatures exceed 90°F.
- Do not graze animals on treated areas.
- Do not feed clippings from treated areas to livestock or poultry.
- Do not apply to apple, Bartlett pear, cherry, citrus, nectarine, peach, pecan, plums or walnut trees that will bear harvestable fruit within 12 months.

MIXING INSTRUCTIONS

- Fill the tank $\frac{1}{2}$ to $\frac{3}{4}$ full of water.
- Add the appropriate amount of LESCO® Spectator™ Turf and Ornamental Fungicide.
- Add the remaining water.
- Thoroughly agitate during mixing and during application.

If this product is mixed with other products, use the following mixing sequence:

1. Check the compatibility of the tank mixture using a jar test before mixing in the spray tank (see Tank Mixes Section).
2. Provide sufficient jet or mechanical agitation during filling and application to keep the tank mix uniformly suspended.
3. Fill tank at least $\frac{1}{2}$ full of water.
4. Add wettable powders to the tank first, allowing them to completely suspend in the water before proceeding. This process can be hastened by premixing the product in water before adding to the tank.
5. Add flowables or suspensions next.
6. Add LESCO® Spectator™ Turf and Ornamental Fungicide.
7. Add other emulsifiable concentrates.
8. Do not leave tank mixtures in the spray tank for prolonged periods without agitation. Mix and apply the same day.

TANK MIXES

LESCO® Spectator™ Turf and Ornamental Fungicide can be tank mixed with other fungicides, herbicides and insecticides. Prior to using the tank mix, determine compatibility of the tank mix components by conducting a jar test:

1. Use a small container, such as a jar, and mix a small amount (0.5 to 1 quart) of spray, combining all ingredients in the same ratio as the anticipated use.
2. Mix thoroughly.
3. After 5 to 15 minutes, observe the mixture for any signs of incompatibility. These can include development of a precipitate, surface oil, or a thick (clabbered) texture.
4. If there are any indications of incompatibility, do not use the tank mix or use Unite compatibility agent (3 pts/100 gallons). Rerun the jar test using Unite at the appropriate ratio to ensure compatibility.

Observe all directions and the most restrictive precautions and limitations on the labels of each product used in the tank mixture.

TURFGRASSES

General Information/Restrictions:

- Do not apply more than 5.8 fl. oz. per 1000 sq. ft.
- Apply in sufficient water to ensure thorough coverage.
- Apply after mowing or allow sprayed areas to dry completely before irrigating.
- Allow sprayed areas to dry completely before irrigating when controlling foliar diseases.
- LESCO® Spectator™ Turf and Ornamental Fungicide can be watered-in immediately after application when controlling soil-borne diseases.
- Under high disease pressure or conditions conducive to high disease pressure, use the higher recommended rate and shorter application interval.
- Prior to use, evaluate spray additives.
- For best control and turf quality, use LESCO® Spectator™ Turf and Ornamental Fungicide in a preventative disease control program and with good turf management practices. Also, properly identify the organism causing the disease.

- Bermudagrass may be sensitive to LESCO® Spectator™ Turf and Ornamental Fungicide. Do not exceed 1.44 fl. oz. per 1000 sq. ft. per 30 days on any variety of bermudagrass.
- In Florida, do not apply LESCO® Spectator™ Turf and Ornamental Fungicide to bermudagrass golf course greens when temperatures exceed 90°F.
- Do not graze animals on treated areas.
- Do not feed clippings from treated areas to livestock or poultry.

Rate Conversion:

Fluid ounces to Milliliters Conversion table				
0.18 fl. oz.	0.36 fl. oz.	0.72 fl. oz.	1.1 fl. oz.	1.44 fl. oz.
5.4 ml	10.8 ml	21.6 ml	33 ml	43.2 ml

1 fl. oz. = 30 milliliters

Disease Control:

Disease	Application Rate		Application Timing or Interval	Instructions
	(fl. oz. per 1000 sq. ft.)	(fl. oz. per acre)		
Anthracnose <i>(Colletotricum graminicola)</i>	0.36-0.72	15.9-31.8	14-28 days	Apply LESCO® Spectator™ Turf and Ornamental Fungicide when conditions are favorable for disease development. Use the higher application rate and shorter application interval when disease pressure is high. Tank mix with a contact fungicide at the labeled rate for broad spectrum control. If disease is present, use the highest rate recommended (i.e. 0.72 fl. oz./1000 sq. ft. or 31.8 fl. oz/A) in combination with a contact fungicide, such as Daconil Ultrex® or Chipco® 26019.

(continued)

Disease	Application Rate		Application Timing or Interval	Instructions
	(fl. oz. per 1000 sq. ft.)	(fl. oz. per acre)		
Brown Patch <i>(Rhizoctonia solani)</i>	0.36-0.72	15.9-31.8	14-21 days	Apply in May or June before disease is present and tank mix with a contact fungicide, such as Daconil Ultrex® or Chipco® 26019, registered for control of brown patch. Use the higher application rate and shorter application interval when disease pressure is high.
Dichondra rust <i>(Puccinia dichondrae)</i>	0.72	31.8	14-21 days	Apply LESCO® Spectator™ Turf and Ornamental Fungicide when conditions are favorable for disease development.

(continued)

Disease	Application Rate		Application Timing or Interval	Instructions
	(fl. oz. per 1000 sq. ft.)	(fl. oz. per acre)		
Dollar spot <i>(Sclerotinia homeocarpa)</i>	0.18	7.9	7 days	Apply LESCO® Spectator™ Turf and Ornamental Fungicide when conditions are favorable for disease development.
	0.18	7.9	14 days	Tank mix with low labeled rate Daconil Ultrex® or Chipco® 26019.
	0.36	15.9	21-28 days	Tank mix with low labeled rate of a product containing chlorothalonil.
	0.36-0.72	15.9-31.8	14-28 days	If LESCO® Spectator™ Turf and Ornamental Fungicide is not tank mixed at this rate, do not make more than 3 consecutive applications for control of dollar spot before rotating to an alternate fungicide with a different mode of action than LESCO® Spectator™ Turf and Ornamental Fungicide.

(continued)

Disease	Application Rate		Application Timing or Interval	Instructions
	(fl. oz. per 1000 sq. ft.)	(fl. oz. per acre)		
Fusarium Patch (<i>Fusarium nivale</i>)	0.72-1.44	31.8-63.4	Fall-Early Spring	Apply LESCO® Spectator™ Turf and Ornamental Fungicide when conditions are favorable for disease development.
Gray Leafspot (<i>Pyricularia grisea</i>)	0.36-0.72	15.9-31.8	14 days	Apply LESCO® Spectator™ Turf and Ornamental Fungicide when conditions are favorable for disease development. If using the lower recommended rate, tank mix with a contact fungicide.
Melting Out, Leaf Spot (<i>Bipolaris</i> spp.) (<i>Drechslera</i> spp.)	0.36-0.72	15.9-31.8	14 days	Tank mix with a contact fungicide for broad spectrum control. To reduce the severity of leaf spot and melting out caused by <i>Helminthosporium</i> -type pathogens and under light to moderate pressure, tank mix using 0.36 fl. oz./1000 sq. ft. LESCO® Spectator™ Turf and Ornamental Fungicide with a contact fungicide.

Disease	Application Rate		Application Timing or Interval	Instructions
	(fl. oz. per 1000 sq. ft.)	(fl. oz. per acre)		
Necrotic Ring spot <i>(Leptosphaeria korrae)</i>	1.44	63.4	Fall or Spring	Apply in fall and/or the early spring. Contact your local extension expert for specific application recommendations.
Pink Patch <i>(Limonomyces roseipellis)</i>	0.72	31.8	14-21 days	Apply LESCO® Spectator™ Turf and Ornamental Fungicide when conditions are favorable for disease development.
Powdery Mildew <i>(Erysiphe graminis)</i>	0.36-0.72	15.9-31.8	14-28 days	Apply LESCO® Spectator™ Turf and Ornamental Fungicide when conditions are favorable for disease development. If disease is present, use the highest recommended rate.
Red Thread <i>(Laetisaria fuciformis)</i>	0.72	31.8	14-21 days	Apply LESCO® Spectator™ Turf and Ornamental Fungicide when conditions are favorable for disease development.

(continued)

Disease	Application Rate		Application Timing or Interval	Instructions
	(fl. oz. per 1000 sq. ft.)	(fl. oz. per acre)		
Rust (<i>Puccinia graminis</i>)	0.36-0.72	15.9-31.8	14-28 days	Apply LESCO® Spectator™ Turf and Ornamental Fungicide when conditions are favorable for disease development. If disease is present, use the highest recommended rate.
Snow Mold Gray (<i>Typhula</i> spp.) Pink (<i>Microdochium nivale</i>)	0.72-1.44	31.8-63.4	Late Fall	Make one application in the late fall before snow cover; do not apply on top of snow. For optimum disease control, tank mix with products containing PCNB or chlorothalonil.
Spring Dead spot (<i>Leptosphaeria korrae</i> , <i>Leptosphaeria narmari</i> , <i>Ophiosphaerella herpotricha</i> , <i>Gaeumannomyces graminis</i>)	1.44	63.4	30 days	Make 1 to 3 applications. If a single application is made, apply in September or October. For multiple applications, begin spraying in August.

Disease	Application Rate		Application Timing or Interval	Instructions
	(fl. oz. per 1000 sq. ft.)	(fl. oz. per acre)		
Stripe smut <i>(Ustilago striiformis)</i> <i>(Urocystis agropyri)</i>	0.36-0.72	15.9-31.8	Fall or Spring	Apply once in the fall after grass becomes dormant or in the early spring before grass starts to grow.
Summer Patch Poa Patch <i>(Magnaporthe poae)</i>	0.72 1.44	31.8 63.4	14 days 28 days	Begin applications in April.
Take-All Patch <i>(Gaeumannomyces graminis)</i>	0.72-1.44	31.8-63.4	Spring and Fall	LESCO® Spectator™ Turf and Ornamental Fungicide reduces the severity of take-all patch. Make 1 or 2 fall applications in September and October or when night temperatures drop below 55°F and 1 or 2 applications in April and May. Contact your local extension expert for specific application recommendations.

(continued)

Disease	Application Rate		Application Timing or Interval	Instructions
	(fl. oz. per 1000 sq. ft.)	(fl. oz. per acre)		
Yellow patch <i>(Rhizoctonia cerealis)</i>	1.1-1.44	48.1-63.4	Late Fall	Make one application in the late fall before snow cover; do not use on top of snow. If using the lower recommended rate, tank mix with a contact fungicide.
Zoysia Patch, Large patch of zoysia <i>(Rhizoctonia solani)</i>	1.1-1.44	48.1-63.4	Early fall	Make one application in the early fall (mid-September to mid-October) prior to development of disease symptoms. Contact your local extension expert for specific application recommendations.

Establishment of Cool Season Turfgrass:

LESCO® Spectator™ Turf and Ornamental Fungicide will improve the establishment rate when it is applied to cool season grass seedlings or sod.

New Seedlings: At the 2 to 3 leaf stage of growth, apply 0.36 fl. oz./1000 sq. ft. for faster root development and top growth.

Sod: Two to six weeks, prior to cutting, apply 0.36 fl. oz./1000 sq. ft. to increase sod knitting and faster establishment after laying.

**ORNAMENTALS, NON-BEARING FRUIT AND NUT TREES
IN NURSERIES AND LANDSCAPING**

General Information/Restrictions:

- Allow spray to dry before overhead irrigation is applied.
- For best control and turf quality, use LESCO® Spectator™ Turf and Ornamental Fungicide in a preventative disease control program and with sound disease management practices.
- Up to 2 gallons of LESCO® Spectator™ Turf and Ornamental Fungicide can be applied per acre/crop/year for outdoor uses.
- For landscape plantings, apply 2.2-2.9 fl. oz./100 gallons of water every 21 days. Begin application before disease development for best control.
- For ornamentals, apply at 0.72-8.7 fl. oz./100 gallons of water (see below for specific recommendations).
- Ornamental plant species not listed on this label may be injured by application of LESCO® Spectator™ Turf and Ornamental Fungicide. Prior to using LESCO® Spectator™ Turf and Ornamental Fungicide on plants not listed on this label, test for phytotoxicity on a small-scale basis.
- LESCO® Spectator™ Turf and Ornamental Fungicide may not be effective against diseases not listed on this label. Prior to using LESCO® Spectator™ Turf and Ornamental Fungicide against diseases not listed on this label, test for efficacy on a small-scale basis.
- Do not apply LESCO® Spectator™ Turf and Ornamental Fungicide to African violets, begonias, Boston fern, or geraniums.
- Prior to large-scale application, evaluate LESCO® Spectator™ Turf and Ornamental Fungicide for phytotoxicity and disease control on a small-scale basis.
- Do not apply to apple, Bartlett pear, cherry, citrus, nectarine, peach, pecan, plums or walnut trees that will bear harvestable fruit within 12 months.

Ornamental Plant Species:

LESCO® Spectator™ Turf and Ornamental Fungicide may be used to control plant diseases on the following plant species (Ornamental plant species not listed on this label may be injured by application of LESCO® Spectator™ Turf and Ornamental Fungicide. Prior to using LESCO® Spectator™ Turf and Ornamental Fungicide on plants not listed on this label, test for phytotoxicity on a small-scale basis.):

Herbaceous Ornamentals

Calendula
Carnation
Chrysanthemum
Delphinium
Gomphrena
Iris
Marigold
Monarda
Phlox
Snapdragon
Sweet William (*Dianthus barbatus*)
Zinnia

Woody Ornamentals

Amelanchier
Ash
Azalea
Crabapple
Crape Myrtle
Dogwood
Douglas Fir
Hawthorn
Juniper
Lilac
Oaks
Pines
Poplars
Pyracantha
Red Tip Photinia
Rhododendron
Roses (outdoor use only)
Shasta Fir

Non-bearing Fruit and Nut Trees (Nursery and Landscape Plantings)

Apple
Bartlett pear
Cherry
Citrus
Nectarine
Peach
Pecan
Plum
Walnut

Please refer to the table below (Ornamental Disease Control) for specific rate and application timing information.

Ornamental Disease Control:

Disease	Ornamental Plant Species	Rate (fl. oz. per 100 gallons)	Application Timing or Interval	Instructions
Conifer Blights				
<i>Phomopsis juniperovora</i> (Phomopsis Blight)	Juniper	1.8-2.9	14-21 days	Apply as a full coverage spray to the point of drip. Begin applying LESCO® Spectator™ Turf and Ornamental Fungicide when junipers start to grow and repeat applications during the period of active growth.
<i>Sirococcus strobolinus</i> (Tip Blight)	Pines	5.8	Mid-late winter and 2 additional applications at 2 month intervals	Apply as a full coverage spray to the point of drip. Make initial application in mid-late winter followed by two additional applications at 2 month intervals.
<i>Sphaeropsis sapinea</i> (Diplodia Tip Blight)	Pines	1.8-2.9	14 days	Apply as a full coverage spray to the point of drip. Prior to the major period of infection, make 3 applications 14 days apart.

Disease	Ornamental Plant Species	Rate (fl. oz. per 100 gallons)	Application Timing or Interval	Instructions
Flower Blights				
<i>Ascochyta chrysanthemi</i> (Ray Blight)	Chrysanthemum	4.3	7 days	Apply as a full coverage spray to the point of drip. Begin applying LESCO® Spectator™ Turf and Ornamental Fungicide when conditions are favorable for disease development.
		7.2	14 days	
<i>Monilinia</i> spp.	Cherry Nectarine Peach Plum	0.72-1.44	Begin applications when there is 5-10% bloom, repeat at 70-100% bloom	Apply as a full coverage spray to the point of drip.
<i>Ovulinia</i> spp.	Azalea Rhododendron	1.8-2.9	As needed	Apply as a full coverage spray to the point of drip. Begin applying LESCO® Spectator™ Turf and Ornamental Fungicide when conditions are favorable for disease development.

(continued)

Disease	Ornamental Plant Species	Rate (fl. oz. per 100 gallons)	Application Timing or Interval	Instructions
Leaf Blights/Spots				
<i>Alternaria</i> spp.	Gomphrena Marigold	1.8-2.9	As needed	Apply as a full coverage spray to the point of drip. Begin applying LESCO® Spectator™ Turf and Ornamental Fungicide when conditions are favorable for disease development.
<i>Cercospora</i> spp. (Brown Leaf spot)	Pecan	4.3	14 days	Apply as a full coverage spray to the point of drip. Begin application at bud break, making a total of 3 applications.
<i>Cladosporium</i> spp. (Scab)	Pecan	4.3	14 days	Apply as a full coverage spray to the point of drip. Begin application at bud break, making a total of 3 applications.
	Crabapple	2.9-4.3	30 days	Apply as a full coverage spray to the point of drip. Begin applying LESCO® Spectator™ Turf and Ornamental Fungicide when conditions are favorable for disease development.

Disease	Ornamental Plant Species	Rate (fl. oz. per 100 gallons)	Application Timing or Interval	Instructions
<i>Coccomyces hiemalis</i>	Cherry	0.72-1.44	14-21 days	Apply as a full coverage spray to the point of drip. Repeat applications during the period of primary application. Tank mix with a contact fungicide if disease is present.
<i>Cristulariella</i> spp. (Zonate Leaf spot)	Pecan	4.3	14 days	Apply as a full coverage spray to the point of drip. Begin application at bud break, making a total of 3 applications.
<i>Diplocarpon rosae</i> (Blackspot)	Roses (outdoor use only)	1.8-2.9	As needed	Apply as a full coverage spray to the point of drip. Begin applying LESCO® Spectator™ Turf and Ornamental Fungicide when conditions are favorable for disease development. Tank mix with a contact fungicide labeled for blackspot.

(continued)

Disease	Ornamental Plant Species	Rate (fl. oz. per 100 gallons)	Application Timing or Interval	Instructions
<i>Discula</i> spp. (Anthracnose)	Dogwood	2.9-4.3	30 days	Apply as a full coverage spray to the point of drip. Begin applying LESCO® Spectator™ Turf and Ornamental Fungicide when conditions are favorable for disease development.
<i>Fabraea maculata</i> (syn. <i>Entomosporium maculata</i>)	Red Tip Photinia	1.8-2.9	As needed	Apply as a full coverage spray to the point of drip. Begin applying LESCO® Spectator™ Turf and Ornamental Fungicide when conditions are favorable for disease development.
<i>Gnomonia leptostyla</i> (Anthracnose)	Walnut	3.1	14-21 days	Apply as a full coverage spray to the point of drip. Begin applying LESCO® Spectator™ Turf and Ornamental Fungicide when conditions are favorable for disease development.

(continued)

Disease	Ornamental Plant Species	Rate (fl. oz. per 100 gallons)	Application Timing or Interval	Instructions
<i>Heterosporium echinulatum</i>	Sweet William (<i>Dianthus barbatus</i>)	1.8-2.9	As needed	Apply as a full coverage spray to the point of drip. Begin applying LESCO® Spectator™Turf and Ornamental Fungicide when conditions are favorable for disease development.
<i>Mycosphaerella caryigena</i> (Downy Spot)	Pecan	4.3	14 days	Apply as a full coverage spray to the point of drip. Begin application at bud break, making a total of 3 applications.
<i>Mycosphaerella fructicola</i> (Greasy Spot)	Citrus	7.2-8.7	Apply within June to August time period	Apply as a full coverage spray to the point of drip.

(continued)

Disease	Ornamental Plant Species	Rate (fl. oz. per 100 gallons)	Application Timing or Interval	Instructions
<i>Septoria</i> spp. (Leaf scorch)	Rhododendron	2.9-4.3	30 days	Apply as a full coverage spray to the point of drip. Begin applying LESCO® Spectator™ Turf and Ornamental Fungicide when conditions are favorable for disease development.
	Pecan	4.3	14 days	Apply as a full coverage spray to the point of drip. Begin application at bud break, making a total of 3 applications.
	Bayberry Wax myrtle	2.9	30 days	Apply as a full coverage spray to the point of drip. Begin applying LESCO® Spectator™ Turf and Ornamental Fungicide when conditions are favorable for disease development.

(continued)

Disease	Ornamental Plant Species	Rate (fl. oz. per 100 gallons)	Application Timing or Interval	Instructions
<i>Spilocaea pyracanthae</i>	Pyracantha	1.8-2.9	As needed	Apply as a full coverage spray to the point of drip. Begin applying LESCO® Spectator™ Turf and Ornamental Fungicide when conditions are favorable for disease development.
<i>Tubakia dryina</i>	Oaks	5.8	14-28 days	Apply as a full coverage spray to the point of drip. Begin applying LESCO® Spectator™ Turf and Ornamental Fungicide when conditions are favorable for disease development.
<i>Venturia inaequalis</i> (Scab)	Crabapple Apple Bartlett pear	0.72-1.44	14-21 days	Apply as a full coverage spray to the point of drip. Repeat applications during the period of primary application. Tank mix with a contact fungicide if disease is present.

(continued)

Disease	Ornamental Plant Species	Rate (fl. oz. per 100 gallons)	Application Timing or Interval	Instructions
Powdery Mildew				
<i>Erysiphe</i> spp.	Calendula	1.8-2.9	30 days	Apply as a full coverage spray to the point of drip. Begin applying LESCO® Spectator™ Turf and Ornamental Fungicide when conditions are favorable for disease development.
	Delphinium Crape Myrtle	1.8-2.9	As needed	Apply as a full coverage spray to the point of drip. Begin applying LESCO® Spectator™ Turf and Ornamental Fungicide when conditions are favorable for disease development.
<i>Microsphaera</i> spp.	Azalea	2.9-4.3	30 days	Apply as a full coverage spray to the point of drip. Begin applying LESCO® Spectator™ Turf and Ornamental Fungicide when conditions are favorable for disease development.

(continued)

Disease	Ornamental Plant Species	Rate (fl. oz. per 100 gallons)	Application Timing or Interval	Instructions
<i>Didium</i> spp.	Monarda Phlox Zinnia Ash Crabapple Dogwood Lilac Bartlett pear	1.8-2.9	As needed	Apply as a full coverage spray to the point of drip. Begin applying LESCO® Spectator™ Turf and Ornamental Fungicide when conditions are favorable for disease development.
<i>Podosphaera</i> spp.	Amelanchier Apple	1.8-2.9	As needed	Apply as a full coverage spray to the point of drip. Begin applying LESCO® Spectator™ Turf and Ornamental Fungicide when conditions are favorable for disease development.
<i>Sphaerotheca pannosa</i>	Roses (outdoor use only) Pecan	1.8-2.9	As needed	Apply as a full coverage spray to the point of drip. Begin applying LESCO® Spectator™ Turf and Ornamental Fungicide when conditions are favorable for disease development.

(continued)

Disease	Ornamental Plant Species	Rate (fl. oz. per 100 gallons)	Application Timing or Interval	Instructions
Rust				
<i>Gymnosporangium juniperi-virginianae</i>	Crabapple Hawthorn Apple Bartlett pear	0.72-1.44	14-21 days	Apply as a full coverage spray to the point of drip. Repeat applications during the period of primary application. Tank mix with a contact fungicide if disease is present.
<i>Melampsora occidentalis</i>	Douglas Fir	5.8	Apply once in May	Apply as a full coverage spray to the point of drip
	Poplars	5.8	14-28 days	Apply as a full coverage spray to the point of drip. Begin applying LESCO® Spectator™ Turf and Ornamental Fungicide when conditions are favorable for disease development.

(continued)

Disease	Ornamental Plant Species	Rate (fl. oz. per 100 gallons)	Application Timing or Interval	Instructions
<i>Phragmidium</i> spp.	Roses (outdoor use only)	1.8-2.9	As needed	Apply as a full coverage spray to the point of drip. Begin applying LESCO® Spectator™ Turf and Ornamental Fungicide when conditions are favorable for disease development.
<i>Puccinia</i> spp.	Iris Snapdragon	1.8-2.9	As needed	Apply as a full coverage spray to the point of drip. Begin applying LESCO® Spectator™ Turf and Ornamental Fungicide when conditions are favorable for disease development.
<i>Pucciniastrum geoppertianum</i>	Shasta Fir	5.8	14-28 days	Apply as a full coverage spray to the point of drip. Begin applying LESCO® Spectator™ Turf and Ornamental Fungicide when conditions are favorable for disease development.

(continued)

Disease	Ornamental Plant Species	Rate (fl. oz. per 100 gallons)	Application Timing or Interval	Instructions
<i>Uromyces dianthi</i>	Carnation	1.8-2.9	As needed	Apply as a full coverage spray to the point of drip. Begin applying LESCO® Spectator™Turf and Ornamental Fungicide when conditions are favorable for disease development.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Store in original container in a secured dry storage area. Prevent cross-contamination with other pesticides and fertilizers.

PESTICIDE DISPOSAL: Pesticide wastes may be acutely hazardous. Improper disposal is a violation of Federal law. Pesticide, mixtures, or equipment rinse water that cannot be chemically reprocessed must be disposed of according to applicable Federal, state or local procedures. Contact your state pesticide or environmental control agency or the hazardous waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

CONDITION OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather, presence of other materials or other influencing factors in the use of the product, which are beyond the control of Etigra or Seller. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold Etigra and Seller harmless for any claims relating to such factors.

Etigra warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. To the extent consistent with applicable law, this warranty does not extend

to the use of this product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller or Etigra, and Buyer and User assume the risk of any such use. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, ETIGRA MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

To the extent allowed by state law, neither Etigra or Seller shall be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF ETIGRA AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF ETIGRA OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

Etigra and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability. This limitation of Warranty and Liability may not be amended by any oral or written agreement.

Daconil Ultrex® is a registered trademark of a Syngenta Group Company.

Chipco® is a registered trademark of Bayer Ag.

Etigra™ is a trademark of Etigra.

LESCO®, Spectator™ and the sweeping design are trademarks of LESCO Technologies, LLC.

EPA 20060817

LESCO
SPECTATOR™
 Turf and Ornamental
 Fungicide

PEEL HERE TO OPEN ▲

#450267

NET CONTENTS: 1 qt

For use on turf and ornamentals and non-bearing fruit and nut trees in nursery and landscape plantings

ACTIVE INGREDIENT:

Propiconazole: 1-[[2-(2,4-dichlorophenyl)-4-propyl-1,3-dioxolan-2-yl]methyl] 41.8%
 -1H-1,2,4-Triazole 58.2%
OTHER INGREDIENTS: 100.0%
TOTAL: 100.0%

Contains petroleum distillates. Contains 4 lbs. of propiconazole per gallon.

**KEEP OUT OF REACH OF CHILDREN
 WARNING AVISO**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand this label, find someone to explain it to you in detail).

See inside label booklet for additional PRECAUTIONARY STATEMENTS

EPA Reg. No. 79676-21 EPA Est. No. indicated by the 8th digit of the batch number on this package. (A) = 4-NY-001; (P) = 34704-MS-002
 Manufactured for:
 Etigra • 501 Cascade Pointe Lane, Suite 103, Cary, NC 27513 • www.etigra.com

PF-31614

PROOF

THIS PROOF IS TO BE CHECKED FOR ACCURACY

Please review and approve **Text, Spelling, Copy Placement, Size, Shape, Colors, Unwind, and Dieline.**

Authorized signature accepts responsibility for accuracy of all copy, color break and artwork. Cimarron Label is not liable for any discrepancies subsequently identified.

PLEASE NOTE: Due to color variance between printers/monitors, the colors represented by this proof cannot be deemed accurate. Please refer to a color matching system such as the Pantone Matching System for a truer representation of spot colors. **THIS PROOF IS NOT ACCURATE FOR COLOR-MATCH.**

WE CANNOT PROCESS THIS ORDER WITHOUT AN AUTHORIZED SIGNATURE


ARTWORK IS APPROVED

REVISED PROOF NEEDED

Signed _____


Date _____

Please Return To: _____



4201 North Westport Avenue, Sioux Falls, South Dakota 57107
 Phone: (605) 978-0451 • Fax: (605) 978-0463

Please indicate the correct unwind position



PROOF DATE: October 24, 2007

CUSTOMER: ETIGRA

JOB NUMBER: 31614

LABEL SIZE: 3.625" x 4"

LABEL COLORS: BLACK

LEAFLET "IN" COLORS: BLACK

LEAFLET "OUT" COLORS: BLACK Orange 021

dieline does not print